

This patient group direction (PGD) must only be used by registered health professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used.

Patient Group Direction (PGD)

For the administration or supply of

Povidone Iodine 5% sterile eye drops

By registered health care professionals for

Use as an Antiseptic

Throughout the Manx Care and those contracted by the Manx Care where appropriate within practice

PGD NUMBER 126

1. Change history

| Version number | Change details | Date |
|----------------|-----------------------|-----------|
| 1 | Original PGD ratified | June 2021 |
| | | |
| | | |

2. Medicines practice guideline 2: *Patient group directions*

Refer to the relevant sections of NICE medicines practice guideline 2: *Patient group directions* as stated in the blank template notes. For further information about PGD signatories, see the NHS and Manx Care [PGD website FAQs](#).

3. PGD development

Refer to [the NICE PGD competency framework for people developing PGDs](#).

| Job Title & organisation | Name | Signature | Date |
|---------------------------------|------|-----------|------|
| Author of the PGD | | | |
| Member of the PGD working group | | | |

4. PGD authorisation

Refer to the [NICE PGD competency framework for people authorising PGDs](#).

| Job Title | Name | Signature | Date |
|--|------|-----------|------|
| Medical Director | | | |
| Chief Pharmacist/ Pharmaceutical Adviser | | | |
| Senior Paramedic | | | |
| Director of Nursing | | | |
| GP Adviser | | | |
| Senior Microbiologist (if PGD contains antimicrobials) | | | |

5. PGD adoption by the provider

Refer to the [NICE PGD competency framework for people authorising PGDs](#).

| Job title and organisation | Signature | Date | Applicable or not applicable to area |
|----------------------------|-----------|------|--------------------------------------|
| | | | |

6. Training and competency of registered healthcare professionals, employed or contracted by the Manx Care, GP practice or Hospice

Refer to the [NICE PGD competency framework for health professionals using PGDs](#)

| | Requirements of registered Healthcare professionals working under the PGD |
|---|---|
| Qualifications and professional registration | <ul style="list-style-type: none"> Registered healthcare professionals, working within or contracted by the Manx Care, GP practice or Hospice who are permitted staff groups outlined within the current PGD policy Pharmacists must be practising in Manx Care authorised premises i.e. contracted pharmacy premises |
| Initial training | <ul style="list-style-type: none"> Knowledge of current guidelines and the administration of the drug specified in this PGD/BNF and of the inclusion and exclusion criteria Training which enables the practitioner to make a clinical assessment to establish the need for the medication covered by this PGD Local training in the use of PGDs |
| Competency assessment | Staff will be assessed on their knowledge of drugs and clinical assessment as part the competency framework for registered health professionals using PGDs |
| Ongoing training and competency | The registered health care professionals should make sure they are aware of any changes to the recommendations for this medication; it is the responsibility of the registered health care professionals to keep up to date with continuing professional development. PGD updates will be held every two years |

7. Clinical Conditions

| | |
|--|--|
| Clinical condition or situation to which this PGD applies | To produce peri-ocular and conjunctival antiseptics prior to intra-vitreous injections |
| Inclusion criteria | Patients over 16 years in preparation for intra-vitreous injections |
| Exclusion criteria | <ul style="list-style-type: none"> • Allergy or hypersensitivity to povidone iodine 5% eye drops • Pregnancy • Breastfeeding |
| Cautions (including any relevant action to be taken) | <ul style="list-style-type: none"> • On instillation an initial burning sensation may be experienced which may last up to 30 seconds • Individuals should be advised not to drive or operate hazardous machinery until normal vision is restored • Systemic absorption may be reduced by compressing the lacrimal sac at the medial canthus for a minute during the following instillation • Side effects are rare but can cause conjunctival hyperaemia and cytotoxicity on deep tissue |
| Arrangements for referral for medical advice | Patient should be referred to a more experienced clinical practitioner for further assessment |
| Action to be taken if patient excluded | Patient should be referred to a more experienced clinical practitioner for further assessment |
| Action to be taken if patient declines treatment | <ul style="list-style-type: none"> • A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment • This information must be documented in the patients' health records • Any patient who declines care must have demonstrated capacity to do so • Where appropriate care should be escalated |

8. Details of the medicine

| | |
|--|---|
| Name, form and strength of medicine | Povidone Iodine 5% eye drops, solution |
| Legal category | Prescription Only Medication (POM) |
| Indicate any off-label use (if relevant) | None |
| Route/method of administration | Topical application to the eyes |
| Dose and frequency | 1 drop per procedure |
| Quantity to be administered and/or supplied | 1 drop about 5 minutes prior to intra-vitreous injection |
| Maximum or minimum treatment period | 2 doses per procedure, ensuring the final drop is administered no longer than 5 minutes prior to intra-vitreous injection |

| | |
|---------------------------|--|
| Storage | <ul style="list-style-type: none"> • Store between 2°C and 8°C • Store in the original package to protect from light • The product may be stored without refrigeration at not more than 25°C for up to one month |
| Adverse effects | <ul style="list-style-type: none"> • In very rare cases, some patients with severe damage to the clear layer at the front of the eye (the cornea) have developed cloudy patches on the cornea due to calcium build-up during treatment • Allergic type reactions, including difficulty breathing or swallowing, swelling of the face, lips, throat or tongue (anaphylactic shock and anaphylactoid reaction), skin rash with irritation and hives - red, raised, itchy bumps (urticaria) • Contact dermatitis. Irritant effect on the membrane that lines the eyelid and the white of the eye causing redness, blisters and itching • Regular and prolonged application of Minims Povidone Iodine may have an effect on the thyroid, causing the thyroid gland to not make enough thyroid hormones to meet the body's needs or in some cases resulting in an overactive thyroid gland, in the elderly • Conjunctival redness – redness of the whites of the eyes • Eye irritation, superficial pinpoint inflammation of the surface of the eye (superficial punctate keratitis and epitheliopathy), dry eye syndrome (keratoconjunctivitis sicca) • Residual yellow discoloration of the eyes and transient brown discoloration of the skin |
| Records to be kept | The administration of any medication given under a PGD must be recorded within the patient's medical records |

9. Patient information

| | |
|---|--|
| Verbal/Written information to be given to patient or carer | <ul style="list-style-type: none"> • On instillation an initial burning sensation may be experienced which may last up to 30 seconds • May cause transient blurring of vision on instillation. Warn patients not to drive or operate hazardous machinery until vision is clear • Verbal information must be given to patients and or carers for all medication being administered under a PGD • Where medication is being supplied under a PGD, written patient information leaflet must also be supplied • A patient information leaflet is available on request |
| Follow-up advice to be given to patient or carer | If symptoms do not improve or worsen or you become unwell, seek medical advice immediately |

10. Appendix A

| References |
|---|
| <ol style="list-style-type: none">1. British National Formulary (BNF) available online: https://bnf.nice.org.uk2. Nursing and Midwifery (2018) “The code” available online: https://www.nmc.org.uk3. Current Health Care Professions Council standards of practice4. General Pharmaceutical Council standards5. The General Optical Council6. Electronic medicines compendium available online: https://www.medicines.org.uk |

11. Appendix B

| Health professionals agreed to practice |
|---|
| <ul style="list-style-type: none">• Each registered healthcare professional will hold their own Competency framework which will be signed and agreed by their mentor• A mentor is defined within the Manx Care policy as any ward/area managers, sisters, senior nurses, GPs, pharmacists or senior paramedics who has completed the PGD training themselves |