

This patient group direction (PGD) must only be used by registered health professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used.

Patient Group Direction (PGD)

For the administration or supply of

Benzylpenicillin Sodium 600mg Intravenous/Intramuscular

By registered health care professionals for

Suspected Meningococcal disease

Throughout the Manx Care and those contracted by the Manx Care where appropriate within practice

PGD NUMBER 128

1. Change history

Version number	Change details	Date
1	Original PGD ratified	June 2021

2. Medicines practice guideline 2: *Patient group directions*

Refer to the relevant sections of NICE medicines practice guideline 2: *Patient group directions* as stated in the blank template notes. For further information about PGD signatories, see the NHS and Manx Care [PGD website FAQs](#)

3. PGD development

Refer to the [NICE PGD competency framework for people developing PGDs](#)

Job Title & organisation	Name	Signature	Date
Author of the PGD			
Member of the PGD working group			

4. PGD authorisation

Refer to the [NICE PGD competency framework for people authorising PGDs](#)

Job Title	Name	Signature	Date
Medical Director			
Chief Pharmacist/ Pharmaceutical Adviser			
Senior Paramedic			
Director of Nursing			
GP Adviser			
Senior Microbiologist (if PGD contains antimicrobials)			

5. PGD adoption by the provider

Refer to the [NICE PGD competency framework for people authorising PGDs](#)

Job title and organisation	Signature	Date	Applicable or not applicable to area

6. Training and competency of registered healthcare professionals, employed or contracted by the Manx Care, GP practice or Hospice

Refer to the [NICE PGD competency framework for health professionals using PGDs](#)

	Requirements of registered Healthcare professionals working under the PGD
Qualifications and professional registration	<ul style="list-style-type: none"> Registered healthcare professionals, working within or contracted by the Manx Care, GP practice or Hospice who are permitted staff groups outlined within the current PGD policy Pharmacists must be practising in Manx Care authorised premises i.e. contracted pharmacy premises
Initial training	<ul style="list-style-type: none"> Knowledge of current guidelines and the administration of the drug specified in this PGD/BNF and of the inclusion and exclusion criteria Training which enables the practitioner to make a clinical assessment to establish the need for the medication covered by this PGD Local training in the use of PGDs
Competency assessment	Staff will be assessed on their knowledge of drugs and clinical assessment as part the competency framework for registered health professionals using PGDs
Ongoing training and competency	The registered health care professionals should make sure they are aware of any changes to the recommendations for this medication; it is the responsibility of the registered health care professionals to keep up to date with continuing professional development. PGD updates will be held every two years

7. Clinical Conditions

Clinical condition or situation to which this PGD applies	Suspected meningococcal disease
Inclusion criteria	Suspected meningococcal disease
Exclusion criteria	Penicillin hypersensitivity (second choice Ceftriaxone) Cross allergy to other beta-lactams such as Cephalosporins should be taken into account
Cautions (including any relevant action to be taken)	<ul style="list-style-type: none"> • Hypersensitivity and anaphylaxis • Serum sickness like reactions • Reduce dose in severe renal impairment (consult product literature) • It should be recognised that any patient with a history of allergy, especially to drugs, is more likely to develop a hypersensitivity after administration and if an allergic reaction occurs the drug should be withdrawn and appropriate treatment given • Delayed absorption from the intramuscular depot may occur in diabetics
Arrangements for referral for medical advice	Patient should be referred to a more experienced clinical practitioner for further assessment
Action to be taken if patient excluded	Patient should be referred to a more experienced clinical practitioner for further assessment
Action to be taken if patient declines treatment	<ul style="list-style-type: none"> • A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment • This information must be documented in the patients' health records • Any patient who declines care must have demonstrated capacity to do so • Where appropriate care should be escalated

8. Details of the medicine

Name, form and strength of medicine	Benzylpenicillin Injection 600mg
Legal category	Prescription Only Medicine (POM)
Indicate any <u>off-label use</u> (if relevant)	None
Route/method of administration	Slow Intravenous injection or Intramuscular injection
Dose and frequency	<p>Children 1-11 months old: 300mg (One half of a 600mg vial):</p> <ul style="list-style-type: none"> • Intravenous: Reconstitute each 600mg vial with 3.6ml of water for injections. Once completely dissolved, inject 2ml (300mg) of the resulting solution intravenously over at least one minute • Intramuscular: Reconstitute each 600mg vial with 1.6ml water for injections. Once completely dissolved inject 1ml (300mg) intramuscularly <p>Children 1-9 years old: 600mg (One 600mg vial):</p> <ul style="list-style-type: none"> • Intravenous: Reconstitute each 600mg vial with 5ml water for injection. Once completely dissolved inject the resulting solution intravenously over at least 2 minutes • Intramuscular: Reconstitute each 600mg vial with 1.6ml water for injection and inject the resulting solution intramuscularly <p>Adults and children 10 and over: 1.2g (Two 600mg vials):</p> <ul style="list-style-type: none"> • Intravenous: Reconstitute each 600mg vial with 5ml of water for injection. Once completely dissolved inject the resulting solution from both vials intravenously over at least 4 minutes • Intramuscular: Reconstitute each 600mg vial with 1.6ml water for injection and inject the resulting solution from each vial intramuscularly at separate sites <p>Longer administration time is particularly important when using doses of 50mg/kg or greater to avoid CNS toxicity</p>
Quantity to be administered and/or supplied	One dose as described above
Maximum or minimum treatment period	One dose only
Storage	Room Temperature (store below 25°C)
Adverse effects	<ul style="list-style-type: none"> • Hypersensitivity reactions include urticarial, fever, joint pains, rashes, angioedema and anaphylaxis • Serum sickness like reactions, haemolytic anaemia and interstitial nephritis, neutropenia, thrombocytopenia, coagulation disorders and central nervous toxicity including convulsions reported. (Especially in high doses or in severe renal impairment) • Diarrhoea and antibiotic associated colitis

Records to be kept	The administration of any medication given under a PGD must be recorded within the patient's medical records
---------------------------	--

9. Patient information

Verbal/Written information to be given to patient or carer	<ul style="list-style-type: none"> • Verbal information must be given to patients and or carers for all medication being administered under a PGD • Where medication is being supplied under a PGD, written patient information leaflet must also be supplied • A patient information leaflet is available on request
Follow-up advice to be given to patient or carer	If symptoms do not improve or worsen or you become unwell, seek medical advice immediately

10. Appendix A

References
<ol style="list-style-type: none"> 1. British national formulary (BNF) available online: https://bnf.nice.org.uk 2. Nursing and Midwifery (2018) "The code" available online: https://www.nmc.org.uk 3. Current Health Care Professions Council standards of practice 4. General Pharmaceutical Council standards 5. The General Optical Council 6. Electronic medicines compendium available online: https://www.medicines.org.uk

11. Appendix B

Health professionals agreed to practice
<ul style="list-style-type: none"> • Each registered healthcare professional will hold their own Competency framework which will be signed and agreed by their mentor • A mentor is defined within the Manx Care policy as any ward/area managers, sisters, senior nurses, GPs, pharmacists or senior paramedics who has completed the PGD training themselves