



Publications approval reference: C1395

Patient Group Direction for Comirnaty® COVID-19 mRNA vaccine

This Patient Group Direction (PGD) is for the administration of Comirnaty® COVID-19 mRNA vaccine to individuals in accordance with the national COVID-19 vaccination programme

This PGD is for the administration of Comirnaty® COVID-19 mRNA vaccine by registered healthcare practitioners identified in [Section 3](#).

The national COVID-19 vaccination programme may also be provided under national protocol or on a patient specific basis (that is by or on the direction of an appropriate independent prescriber). Supply and administration in these instances are not covered by this PGD.

Reference no: Comirnaty COVID-19 mRNA vaccine PGD
Version no: v02.00
Valid from: 20 September 2021
Review date: 1 October 2021
Expiry date: 31 March 2022

Public Health England (PHE) has developed this PGD for authorisation by NHS England and NHS Improvement to facilitate the delivery of the national COVID-19 vaccination programme.

NHS England and NHS Improvement and those providing services in accordance with this PGD must not alter, amend or add to the clinical content of this document (sections 3, 4, 5 and 6); such action will invalidate the clinical sign-off with which it is provided. [Section 2](#) may be amended only by the person(s) authorising the PGD, in accordance with Human Medicines Regulations 2012 (HMR2012)¹ [Schedule 16 Part 2](#), on behalf of NHS England and NHS Improvement. [Section 7](#) is to be completed by registered practitioners providing the service and their authorising/line manager.

Operation of this PGD is the responsibility of NHS England and NHS Improvement and service providers. The final authorised copy of this PGD should be kept by NHS England and NHS Improvement for 25 years after the PGD expires. Provider organisations adopting authorised versions of this PGD should also retain copies for the period specified above.

Individual registered practitioners must be authorised by name to work according to the current version of this PGD by signing section 7. A manager with the relevant level of authority should also provide a counter signature, unless there are contractual arrangements for self-declaration.

Providers must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Current versions of PHE developed COVID-19 vaccine PGDs can be found via: [COVID-19 vaccination programme - GOV.UK \(www.gov.uk\)](https://www.gov.uk/government/collections/covid-19-vaccination-programme)

The most current national recommendations should be followed. This may mean that a Patient Specific Direction (PSD) is required to administer the vaccine in line with updated recommendations that are outside the criteria specified in this PGD. Any concerns regarding the content of this PGD should be addressed to: immunisation@phe.gov.uk




¹ This includes any relevant amendments to legislation (such as [2013 No.235](#), [2015 No.178](#), [2015 No.323](#) and [2020 No.1125](#)).

Change history

Version	Change details	Date
V01.00	New PHE PGD template for Comirnaty® COVID-19 mRNA vaccine	06/08/2021
V02.00	<p>PHE PGD template for Comirnaty® COVID-19 mRNA Vaccine V01.00 updated to:</p> <ul style="list-style-type: none"> • remove specific reference to clinically extremely vulnerable (CEV) individuals as they are covered by the inclusion of those in at risk groups • include individuals aged 12 years to under 16 years of age who are in an at-risk group (see the table 'Clinical risk groups for children aged 12-15 years' in Chapter 14a) • include other individuals from age 12 years to under 18 years of age, who do not meet any of the other criteria for inclusion, as eligible for their first dose of the COVID-19 vaccine only • include individuals referred for a third primary dose of COVID-19 vaccine in accordance with patient specific recommendations from their specialist, GP or prescriber • include individuals eligible for a booster dose as part of the national COVID-19 vaccination programme • exclude individuals who have experienced myocarditis or pericarditis determined as likely to be related to previous COVID-19 vaccination • move cautions relating to pregnancy and those involved in clinical trials to the additional information section • update to cautions • update the additional information on immunosuppressed individuals, co-administration and incomplete vaccination • remove key references to JCVI statements which are now incorporated into the guidance in Chapter 14a • minor wording changes and additions to text for consistency; updated references 	15/09/2021

1. PGD development

This PGD has been developed by the following health professionals on behalf of Public Health England:

Developed by:	Name	Signature	Date
Pharmacist (Lead Author)	Elizabeth Graham Lead Pharmacist Immunisation Services, Immunisation and Countermeasures, PHE		18/09/2021
Doctor	Mary Ramsay Consultant Epidemiologist and Head of Immunisation and Countermeasures, PHE		18/09/2021
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant, Immunisation and Countermeasures, PHE		18/09/2021

In addition to the signatories above the working group included:

Name	Designation
Jane Horsfall	Senior Policy Manager, Primary Care Group, NHS England and NHS Improvement
Jo Jenkins	Specialist Pharmacist (Patient Group Directions), NHS Specialist Pharmacy Service
Jill Loader	Deputy Director, Primary Care Group, NHS England and NHS Improvement
Bhavana Reddy	Director of Pharmacy - Clinical Workstream, Flu and COVID-19 Vaccination Programme, NHS England and NHS Improvement
Gul Root	Principal Pharmaceutical Officer, Department of Health and Social Care and National lead pharmacy public health, Public Health England

This PGD has been peer reviewed by the PHE Immunisations PGD Expert Panel in accordance with PHE PGD Policy. It has been ratified by the PHE Medicines Governance Group and the PHE Quality and Clinical Governance Delivery Board.

Expert Panel

Name	Designation
Nicholas Aigbogun	Consultant in Communicable Disease Control, Yorkshire and Humber Health Protection Team, Public Health England
Sarah Dermont	Clinical Project Coordinator and Registered Midwife, NHS Infectious Diseases in Pregnancy Screening Programme, Public Health England
Ed Gardner	Advanced Paramedic Practitioner/Emergency Care Practitioner, Medicines Manager, Proactive Care Lead
Michelle Jones	Senior Medicines Optimisation Pharmacist, NHS Bristol North Somerset and South Gloucestershire CCG
Jacqueline Lamberty	Lead Pharmacist Medicines Management Services, Public Health England
Vanessa MacGregor	Consultant in Communicable Disease Control, Public Health England, East Midlands Health Protection Team


Alison Mackenzie	Consultant in Public Health Medicine, Screening and Immunisation Lead, Public Health England (South West) / NHS England and NHS Improvement South (South West)
Gill Marsh	Principal Screening and Immunisation Manager, Public Health England / NHS England and NHS Improvement (North West)
Lesley McFarlane	Screening and Immunisation Manager: Clinical (COVID-19 and Influenza), Public Health England / NHS England and NHS Improvement (Midlands)
Tushar Shah	Lead Pharmacy Advisor, NHS England and NHS Improvement (London Region)

2. Organisational authorisation

The PGD is not legally valid until it has had the relevant organisational authorisation from NHS England and NHS Improvement completed below.

NHS England and NHS Improvement accepts governance responsibility for this PGD. Any provider delivering the national COVID-19 vaccination programme under PGD must work strictly within the terms of this PGD, relevant NHS standard operating procedures (SOPs) and contractual arrangements with the commissioner for the delivery of the national COVID-19 vaccination programme.

NHS England and NHS Improvement authorises this PGD for use by the services or providers delivering the national COVID-19 vaccination programme.

Organisational approval (legal requirement)			
Role	Name	Sign	Date
Medical Director, COVID-19 Vaccination Programme, NHS England and NHS Improvement	Dr Jonathan Leach OBE		20/09/2021

[Section 7](#) provides a practitioner authorisation sheet. Individual practitioners must be authorised by name to work to this PGD. Alternative practitioner authorisation records, specifying the PGD and version number, may be used where appropriate in accordance with local policy. This may include the use of electronic records.

Assembly, final preparation and administration of vaccines supplied and administered under this PGD must be subject to NHS governance arrangements and standard operating procedures that ensure that the safety, quality or efficacy of the product is not compromised. The assembly, final preparation and administration of the vaccines should also be in accordance with the manufacturer's instructions in the product's UK Summary of Product Characteristics ([SPC](#)) and/or in accordance with official national recommendations.

3. Characteristics of staff

<p>Qualifications and professional registration</p>	<p>Practitioners must only work under this PGD where they are competent to do so. Practitioners working to this PGD must also be one of the following registered professionals who can legally supply and administer under a PGD (see Patient Group Directions: who can administer them):</p> <ul style="list-style-type: none"> • nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) • pharmacists currently registered with the General Pharmaceutical Council (GPhC) • chiropodists/podiatrists, dieticians, occupational therapists, orthoptists, orthotists/prosthetists, paramedics, physiotherapists, radiographers and speech and language therapists currently registered with the Health and Care Professions Council (HCPC) • dental hygienists and dental therapists registered with the General Dental Council • optometrists registered with the General Optical Council. <p>Practitioners must also fulfil all of the Additional requirements.</p>
<p>Additional requirements</p>	<p>Additionally, practitioners:</p> <ul style="list-style-type: none"> • must be authorised by name as an approved practitioner under the current terms of this PGD before working to it • must have undertaken appropriate training for working under PGDs for supply/administration of medicines • must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs) • must be familiar with the vaccine product and alert to changes in the SPC, and familiar with the national recommendations for the use of this vaccine • must be familiar with, and alert to changes in relevant chapters of Immunisation Against Infectious Disease: the Green Book • must be familiar with, and alert to changes in the relevant NHS standard operating procedures (SOPs) and commissioning arrangements for the national COVID-19 vaccination programme • must have undertaken training appropriate to this PGD as required by local policy and SOPs and in line with the Training recommendations for COVID-19 vaccinators. • must have undertaken training to meet the minimum standards in relation to vaccinating those under 18 as required by national and local policy. • must have completed the national COVID-19 vaccination e-learning programme, including the relevant vaccine specific session, and/or locally-provided COVID-19 vaccine training • must be competent to assess individuals for suitability for vaccination, identify any contraindications or precautions, obtain informed consent (or 'best interests' decision in accordance with the Mental Capacity Act 2005) and to discuss issues related to vaccination. For further information on consent see Chapter 2 of 'The Green Book'. • must be competent in the correct handling and storage of vaccines, and management of the cold chain • must be competent in the handling of the vaccine product, procedure for dilution of the vaccine and use of the correct technique for drawing up the correct dose • must be competent in the intramuscular injection technique

Continued over page

<p>Additional requirements (continued)</p>	<ul style="list-style-type: none"> • must be competent in the recognition and management of anaphylaxis, have completed basic life support training and be able to respond appropriately to immediate adverse reactions • must have access to the PGD and relevant COVID-19 vaccination programme online resources such as the Green Book and PHE COVID-19 vaccination programme: Information for healthcare practitioners • must have been signed off as competent using the COVID-19 vaccinator competency assessment tool if new to or returning to immunisation after a prolonged period (more than 12 months) or have used the tool for self-assessment if experienced vaccinator (vaccinated within past 12 months) • should fulfil any additional requirements defined by local or national policy <p>The individual practitioner must be authorised by name, under the current version of this PGD before working according to it.</p>
<p>Continued training requirements</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to vaccination and management of anaphylaxis.</p> <p>Practitioners should be constantly alert to any subsequent recommendations from Public Health England and/or NHS England and NHS Improvement and other sources of medicines information.</p>

4. Clinical condition or situation to which this PGD applies

<p>Clinical condition or situation to which this PGD applies</p>	<p>Comirnaty® COVID-19 mRNA vaccine is indicated for the active immunisation of individuals for the prevention of coronavirus disease (COVID-19) caused by the SARS-CoV-2 virus, in accordance with the national COVID-19 vaccination programme (see COVID-19 vaccination programme page) and recommendations given in Chapter 14a of the Immunisation Against Infectious Disease: the ‘Green Book’, and subsequent correspondence/publications from PHE and/or NHS England and NHS Improvement.</p>
<p>Criteria for inclusion</p>	<p>Comirnaty® COVID-19 mRNA vaccine should be offered in accordance with the national COVID-19 vaccination programme to:</p> <ul style="list-style-type: none"> • all individuals aged 18 years and over • individuals 16 years to 18 years of age who are: <ul style="list-style-type: none"> ○ frontline health and social care workers (see Chapter 14a) ○ in an at-risk group (see the table ‘Clinical and other risk groups 16 years of age and over who should receive primary COVID-19 immunisation’ in Chapter 14a)² • individuals aged 12 years to under 16 years of age who are in an at-risk group (see the table ‘Clinical risk groups for children aged 12-15 years’ in Chapter 14a) • individuals aged 12 years and above who expect to share living accommodation on most days (and therefore for whom continuing close contact is unavoidable) with individuals of any age who are immunosuppressed • other individuals from age 12 years to under 18 years of age who do not meet any of the above criteria for inclusion. This group is currently only eligible for their first dose of the COVID-19 vaccine. A decision on when to offer the second dose to healthy children and young people is pending further evidence (see Dose and frequency of administration section). • individuals referred for a third primary dose of COVID-19 vaccine in accordance with patient specific recommendations from their specialist, GP or prescriber, who has assessed the individual’s indication for a third primary dose of COVID-19 vaccine in accordance with current recommendations from the JCVI and Chapter 14a of the Green Book. Note: The decision on the timing of the third dose should be undertaken by the specialist involved in the care of the patient. • individuals eligible for a booster dose as part of the national COVID-19 vaccination programme in accordance with the recommendations from the JCVI and Chapter 14a of the Green Book.
<p>Criteria for exclusion³</p> <p>Continued over page</p>	<p>Individuals for whom valid consent, or ‘best-interests’ decision in accordance with the Mental Capacity Act 2005, has not been obtained (for further information on consent see Chapter 2 of ‘The Green Book’). The Patient Information Leaflet (PIL) for Comirnaty® COVID-19 mRNA vaccine should be available to inform consent.</p> <p>Individuals who:</p> <ul style="list-style-type: none"> • are less than 12 years of age • have had a previous systemic allergic reaction (including immediate onset anaphylaxis) to a previous dose of a COVID-19 mRNA vaccine

² This also includes adult carers.

³ Exclusion under this PGD does not necessarily mean the medication is contraindicated, but it would be outside its remit and another form of authorisation will be required

<p>Criteria for exclusion (continued)</p>	<p>or to any component of the Comirnaty® COVID-19 mRNA vaccine or residues from the manufacturing process⁴</p> <ul style="list-style-type: none"> • have a history of prior allergic reaction to COVID-19 vaccine that required medical attention • have a history of immediate anaphylaxis to multiple, different drug classes, with the trigger unidentified (this may indicate polyethylene glycol (PEG) allergy) • have a history of anaphylaxis to a vaccine, injected antibody preparation or a medicine likely to contain PEG (such as depot steroid injection, laxative) • have history of idiopathic anaphylaxis • have experienced myocarditis or pericarditis determined as likely to be related to previous COVID-19 vaccination • are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for vaccination) • have received a full dose of COVID-19 vaccine in the preceding 21 days
<p>Cautions, including any relevant action to be taken</p> <p>Continued over page</p>	<p>All recipients of the Comirnaty® COVID-19 mRNA vaccine should be kept for observation and monitored for a minimum of 15 minutes. Facilities for management of anaphylaxis should be available at all vaccination sites.</p> <p>Where individuals experienced a possible allergic reaction to a dose of COVID-19 vaccine, follow the guidance in Chapter 14a of the Green Book in relation to the administration of subsequent doses.</p> <p>Individuals with non-allergic reactions (vasovagal episodes, non-urticarial skin reaction or non-specific symptoms) to a COVID-19 vaccine can receive subsequent doses of vaccine in any vaccination setting.</p> <p>Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.</p> <p>Individuals with a bleeding disorder may develop a haematoma at the injection site. Individuals with bleeding disorders may be vaccinated intramuscularly if, in the opinion of a doctor familiar with the individual's bleeding risk, vaccines or similar small volume intramuscular injections can be administered with reasonable safety by this route. If the individual receives medication/treatment to reduce bleeding, for example treatment for haemophilia, intramuscular vaccination can be scheduled shortly after such medication/treatment is administered. Individuals on stable anticoagulation therapy, including individuals on warfarin who are up to date with their scheduled INR testing and whose latest INR was below the upper threshold of their therapeutic range, can receive intramuscular vaccination. A fine needle (equal to 23 gauge or finer calibre such as 25 gauge) should be used for the vaccination, followed by firm pressure applied to the site (without rubbing) for at least 2 minutes. If in any doubt, consult with the clinician responsible for prescribing or monitoring the individual's anticoagulant therapy. The individual/carer should be informed about the risk of haematoma from the injection.</p> <p>Very rare reports have been received of Guillain-Barre Syndrome (GBS) following COVID-19 vaccination, although no causal link with COVID-19 vaccination has been proven (further information is available in Chapter 14a). Healthcare professionals should be alert to the signs and symptoms of GBS to ensure correct diagnosis and to rule out other</p>

⁴ Contains polyethylene glycol (PEG), refer to the [SPC](#) for a full list of excipients.

<p>Cautions, including any relevant action to be taken (continued)</p>	<p>causes, in order to initiate adequate supportive care and treatment. Individuals who have a history of GBS should be vaccinated as recommended for their age and underlying risk status.</p> <p>Past history of COVID-19 infection</p> <p>There is no evidence of any safety concerns from vaccinating individuals with a past history of COVID-19 infection, or with detectable COVID-19 antibody.</p> <p>Vaccination of individuals who may be infected but asymptomatic or incubating COVID-19 infection is unlikely to have a detrimental effect on the illness. Vaccination should be deferred in those with confirmed infection to avoid onward transmission and confusing the differential diagnosis. As clinical deterioration can occur up to two weeks after infection, ideally vaccination should be deferred until clinical recovery to around four weeks after onset of symptoms or four weeks from the first confirmed positive specimen in those who are asymptomatic.</p> <p>Having prolonged COVID-19 symptoms is not a contraindication to receiving COVID-19 vaccine but if the individual is seriously debilitated, still under active investigation, or has evidence of recent deterioration, deferral of vaccination may be considered to avoid incorrect attribution of any change in the person's underlying condition to the vaccine.</p> <p>Vaccine Surveillance</p> <p>The UK regulator will maintain real-time surveillance post deployment of COVID-19 vaccines in the UK. In response to any safety signals, MHRA may provide temporary advice or make substantive amendments to the authorised conditions of the vaccine product's supply in the UK. Administration under this PGD must be in accordance with the most up-to-date advice or amendments (see Green Book Chapter 14a and the SPC). These documents take precedence for the purposes of compliance with this PGD, if there is a delay in updating other provisions of this PGD that cut across them.</p>
<p>Action to be taken if the patient is excluded</p> <p>Continued over page</p>	<p>The risk to the individual of not being immunised must be considered. The indications for risk groups are not exhaustive, and the healthcare practitioner should consider the risk of COVID-19 exacerbating any underlying disease that an individual may have, as well as the risk of serious illness from COVID-19 itself. Where appropriate, such individuals should be referred for assessment of clinical risk. Where risk is identified as equivalent to those currently eligible for immunisation, vaccination may be provided by an appropriate prescriber or on a patient specific basis, under a PSD.</p> <p>For individuals who have had a previous systemic allergic reaction (including immediate onset anaphylaxis) to a previous dose of COVID-19 mRNA vaccine, or any component of the vaccine, advice should be sought from an allergy specialist.</p> <p>Special precautions as described in Chapter 14a, and consideration of the possibility of undiagnosed PEG-allergy, is required for individuals with:</p> <ul style="list-style-type: none"> • history of prior allergic reaction to COVID-19 vaccine that required medical attention • history of immediate anaphylaxis to multiple, different drug classes, with the trigger unidentified (this may indicate PEG allergy) • history of anaphylaxis to a vaccine, injected antibody preparation or a medicine likely to contain PEG (such as depot steroid injection, laxative) • history of idiopathic anaphylaxis <p>Such individuals should not be vaccinated with Comirnaty® COVID-19</p>

<p>Action to be taken if the patient is excluded (continued)</p>	<p>mRNA vaccine, except on the expert advice of an allergy specialist and under a PSD.</p> <p>Individuals who have experienced myocarditis or pericarditis following COVID-19 vaccination should be assessed by an appropriate clinician to determine whether it is likely to be vaccine related. As the mechanism of action and risk of recurrence of myocarditis and pericarditis are being investigated, the current advice is that an individual's second or subsequent doses should be deferred until further information becomes available; an alternate vaccine may not be safer and will increase the risk of other adverse events.</p> <p>In case of postponement due to acute illness, advise when the individual can be vaccinated and if possible, ensure another appointment is arranged.</p> <p>Document the reason for exclusion and any action taken.</p>
<p>Action to be taken if the patient or carer declines treatment</p>	<p>Informed consent, from the individual or a person legally able to act on the person's behalf, must be obtained for each administration and recorded appropriately. Where a person lacks the capacity, in accordance with the Mental Capacity Act 2005, a decision to vaccinate may be made in the individual's best interests. For further information on consent see Chapter 2 of 'The Green Book'.</p> <p>Advise the individual/carer about the protective effects of the vaccine, the risks of infection and potential complications if not immunised.</p> <p>Document advice given and the decision reached.</p>
<p>Arrangements for referral</p>	<p>As per local policy.</p>

5. Description of treatment

Name, strength and formulation of drug	<p>Comirnaty® concentrate for dispersion for injection COVID-19 mRNA vaccine (nucleoside modified)</p> <p>1 vial (0.45ml) contains 6 doses of 0.3ml after dilution.</p> <p>1 dose (0.3ml) contains 30micrograms of COVID-19 mRNA vaccine (embedded in lipid nanoparticles).</p> <p>Note: Where appropriate to the delivery model, this PGD may also be used for the administration of vaccine that has been prepared (diluted) by another person in accordance with the manufacturer's instructions and Human Medicines Regulation 3A (UK Statutory Instrument 2020 No. 1594), that is prepared by or under the supervision of a doctor, a registered nurse or a pharmacist.</p>
Legal category	<p>Prescription only medicine (POM).</p>
Black triangle▼	<p>Yes. As a new vaccine product, MHRA has a specific interest in the reporting of adverse drug reactions for this product.</p>
Off-label use	<p>The Comirnaty® COVID-19 mRNA vaccine SPC recommends the second dose is administered 3 weeks after the first dose. There is evidence of better immune response and/or protection from COVID-19 vaccines where longer intervals between doses are used. Therefore, Comirnaty® COVID-19 mRNA vaccine should be administered under this PGD in accordance with official national recommendations from the JCVI for the delivery of the COVID-19 vaccination programme in England (see Chapter 14a). At the time of writing, JCVI recommends an interval of 8 weeks between primary doses.</p> <p>The Comirnaty® COVID-19 mRNA vaccine SPC recommends a two dose course. JCVI, as detailed in Chapter 14a of the Green Book, have made specific recommendations regarding individuals who may benefit from a third primary dose. Individuals may be vaccinated under this PGD where they have been referred for a third primary dose of COVID-19 vaccine in accordance with patient specific recommendations from their specialist, GP or prescriber, who has assessed the individual's indication for a third dose of COVID-19 vaccine in accordance with current recommendations from the JCVI and Chapter 14a of the Green Book. Note: Patient specific advice, in relation to eligibility, the appropriate COVID-19 vaccine and timing of the third vaccine dose, should be provided by the specialist, GP or prescriber to inform the vaccine provider.</p> <p>JCVI have also made recommendations regarding individuals who may benefit from a booster dose. Individuals may be vaccinated under this PGD where they are eligible for a booster dose as part of the national COVID-19 vaccination programme in accordance with the recommendations from the JCVI and Chapter 14a of the Green Book.</p> <p>Vaccine should be stored according to the conditions detailed in the Storage section below. However, in the event of an inadvertent or unavoidable deviation of these conditions refer to PHE Vaccine Incident Guidance. Where vaccine is assessed in accordance with these guidelines as appropriate for continued use this would constitute off-label administration under this PGD.</p> <p>Where a vaccine is recommended off-label consider, as part of the consent process, informing the individual/carer that the vaccine is being offered in accordance with national guidance but that this is outside the product licence.</p>

<p>Route / method of administration</p>	<p>Comirnaty® COVID-19 mRNA vaccine is for administration by intramuscular injection only, preferably into deltoid region of the upper arm.</p> <p>Comirnaty® COVID-19 mRNA vaccine requires dilution in its original vial with 1.8ml of unpreserved sodium chloride 0.9% solution for injection, prior to withdrawing a 0.3ml dose for administration.</p> <p>Vaccine should be prepared in accordance with manufacturer's recommendations (see the product's SPC) and NHS standard operating procedures for the service.</p> <p>Frozen vials should be transferred to an environment of 2°C to 8°C to thaw; a 195 vial pack may take 3 hours to thaw.</p> <p>Alternatively, frozen vials may also be thawed for 30 minutes at temperatures up to 30 °C for immediate use.</p> <p>Allow the thawed vial to come to room temperature and gently invert it 10 times prior to dilution. Do not shake.</p> <p>Prior to dilution, the thawed dispersion may contain white to off-white opaque amorphous particles.</p> <p>The thawed vaccine must be diluted in its original vial with 1.8 mL sodium chloride 0.9% solution for injection, using a 21 gauge or narrower needle and aseptic techniques.</p> <p>Equalise vial pressure before removing the needle from the vial stopper by withdrawing 1.8 mL air into the empty diluent syringe.</p> <p>Gently invert the diluted dispersion 10 times. Do not shake the vaccine.</p> <p>The diluted vaccine should present as an off-white dispersion with no particulates visible. Do not use the diluted vaccine if particulates or discolouration are present.</p> <p>The diluted vials should be marked with the appropriate date and time.</p> <p>After dilution store at 2°C to 30 °C and use within 6 hours, including any transportation time.</p> <p>Do not freeze or shake the diluted dispersion. If refrigerated, allow the diluted dispersion to come to room temperature prior to use.</p> <p>The vaccine dose should be drawn up from the diluted vial immediately prior to administration.</p> <p>In order to extract at least 6 doses from a single vial, low dead-volume syringes and/or needles should be used. Each dose must contain 0.3ml of vaccine. If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 ml, discard the vial and any excess volume. Do not pool excess vaccine from multiple vials.</p> <p>Discard any unused vaccine within 6 hours after dilution.</p> <p>Check product name, batch number and expiry date prior to administration.</p>
<p>Dose and frequency of administration</p> <p>Continued over page</p>	<p>Primary vaccination</p> <p>Individuals from 12 years to under 18 years of age who do not meet any of the other criteria for inclusion should receive their first dose only. A decision on when to offer the second dose to healthy young people, under 18 years of age, is pending further evidence on the safety of a second dose in this age group. Individuals who attain 18 years of age are eligible to complete a two-dose course and should be offered a second dose.</p>

<p>Dose and frequency of administration (continued)</p>	<p>A two-dose course should be administered to eligible individuals, with the exception of:</p> <ul style="list-style-type: none"> • individuals from 12 years to under 18 years of age who do not meet any of the other criteria for inclusion, who should receive their first dose only • individuals who were severely immunosuppressed when they received their first or second dose of COVID- 19 vaccination for whom JCVI have provided recommendations for a third primary dose <p>The two-dose course consists of 30micrograms in 0.3ml followed by a second dose of 30micrograms in 0.3ml after an interval of at least 21 days. However, the schedule should usually be administered in accordance with official national guidance which, at the time of writing, recommends a minimum interval of eight weeks between primary doses.</p> <p>There is evidence of better immune response and/or protection where longer intervals between doses in the primary schedule are used. Based on this evidence, longer intervals are likely to provide more durable protection.</p> <p>At the time of writing, JCVI is currently recommending a minimum interval of eight weeks between doses of all the available COVID-19 vaccines where a two-dose primary schedule is used. Operationally, this consistent interval should be used for all vaccines with a two-dose primary schedule to avoid confusion and simplify booking, and this will help to ensure a good balance between achieving rapid and long-lasting protection.</p> <p>If an interval longer than the recommended interval is left between doses, the second dose should still be given (using the same vaccine as was given for the first dose if possible, see Additional Information). The course does not need to be restarted.</p> <p>The main exception to the eight-week lower interval would be those about to commence immunosuppressive treatment. In these individuals, the licensed minimal interval of at least 21 days may be followed to enable the vaccine to be given whilst their immune system is better able to respond.</p> <p>JCVI advises that a third primary dose be offered to individuals aged 12 years and over who had severe immunosuppression in proximity to their first or second COVID-19 doses in the primary schedule (see 'Box: Criteria for a third primary dose of COVID-19 vaccine' in Chapter 14a). The decision on the timing of the third dose should be undertaken by the specialist involved in the care of the individual. The third dose should be given ideally at least 8 weeks after the second dose, with special attention paid to current or planned immunosuppressive therapies (see Additional information).</p> <p>Booster vaccination</p> <p>A booster dose should be offered no earlier than six months after completion of the primary course, of an mRNA or adenovirus-vectored COVID-19 vaccine, to individuals eligible for a booster (third) dose as part of the national COVID-19 vaccination programme in accordance with the recommendations from the JCVI and Chapter 14a of the Green Book.</p>
<p>Duration of treatment</p>	<p>See Dose and frequency of administration above.</p>
<p>Quantity to be supplied / administered</p>	<p>Administer 30micrograms in 0.3ml per dose.</p>

<p>Supplies</p>	<p>Providers should order/receive COVID-19 vaccines via the national appointed supply route for the provider.</p> <p>NHS standard operating procedures should be followed for appropriate ordering, storage, handling, preparation, administration and waste minimisation of Comirnaty® COVID-19 mRNA Vaccine, which ensure use is in accordance with product's SPC and official national recommendations.</p>
<p>Storage</p>	<p>Comirnaty® COVID-19 mRNA Vaccine is supplied from the manufacturer as a multiple-dose vial of frozen, preservative-free concentrate, which requires storage at -90°C to -60°C.</p> <p>Frozen Vial</p> <p>Shelf life is 6 months at -90°C to -60°C</p> <p>Within the 6 months shelf life, unopened vials may be stored and transported at -25°C to -15°C for a single period of up to 2 weeks and can be returned to -90°C to -60°C.</p> <p>Thawed vial</p> <p>Thawed unopened vials have a 1-month shelf-life at 2°C to 8°C.</p> <p>Within the 1-month shelf-life at 2°C to 8°C, up to 12 hours may be used for transportation.</p> <p>Prior to use, the unopened vaccine can be stored for up to 2 hours at temperatures up to 30°C.</p> <p>Store in original packaging in order to protect from light. During storage, minimise exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. Thawed vials can be handled in room light conditions.</p> <p>Once a vial is removed from the tray, it should be thawed for use.</p> <p>Once thawed the vaccine cannot be re-frozen.</p> <p>Diluted product</p> <p>Chemical and physical in-use stability, including during transportation, has been demonstrated for 6 hours at 2°C to 30°C after dilution in sodium chloride 0.9% solution for injection. From a microbiological point of view, unless the method of dilution precludes the risk of microbial contamination, the product should be used immediately.</p> <p>Precautions for storage</p> <p>Store in original packaging in order to protect from light.</p> <p>During storage, minimise exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.</p> <p>Thawed vials can be handled in room light conditions.</p> <p>These details relate to storage requirements and available stability data at the time of product authorisation. This may be subject to amendment as more data becomes available. Refer to NHS standard operating procedures for the service and the most up to date manufacturer's recommendations in the product's SPC. The product's SPC also contains further information on stability to guide healthcare professionals only in case of temporary temperature excursion.</p> <p>In the event of an inadvertent or unavoidable deviation of these conditions, vaccine that has been stored outside the conditions stated above should be quarantined and risk assessed for suitability of</p>

Continued over page

Storage (continued)	continued off-label use or appropriate disposal. Refer to PHE Vaccine Incident Guidance .
Disposal	<p>Follow local clinical waste policy and NHS standard operating procedures and ensure safe and secure waste disposal.</p> <p>Equipment used for vaccination, including used vials, ampoules, or discharged vaccines in a syringe or applicator, should be disposed of safely and securely according to local authority arrangements and guidance in the technical memorandum 07-01: Safe management of healthcare waste (Department of Health, 2013).</p>
Drug interactions	<p>Immunological response may be diminished in those receiving immunosuppressive treatment, but it is important to still immunise this group.</p> <p>Although no data for co-administration of COVID-19 vaccine with other vaccines exists, in the absence of such data, first principles would suggest that interference between inactivated vaccines with different antigenic content is likely to be limited. Based on experience with other vaccines, any potential interference is most likely to result in a slightly attenuated immune response to one of the vaccines. There is no evidence of any safety concerns, although it may make the attribution of any adverse events more difficult. Similar considerations apply to co-administration of inactivated (or non-replicating) COVID-19 vaccines with live vaccines such as MMR. In particular, live vaccines which replicate in the mucosa, such as live attenuated influenza vaccine (LAIV) are unlikely to be seriously affected by concomitant COVID-19 vaccination.</p> <p>For further information about co-administration with other vaccines see Additional Information section.</p>
Identification and management of adverse reactions Continued over page	<p>The most frequent adverse reactions in individuals 16 years of age and older are injection site pain, fatigue, headache, myalgia, chills, arthralgia, pyrexia and injection site swelling. These reactions are usually mild or moderate in intensity and resolve within a few days after vaccination. Redness at the injection site, nausea and vomiting are reported as common. Lymphadenopathy is reported with a frequency of less than 1%.</p> <p>The most frequent adverse reactions in individuals 12 to 15 years of age are injection site pain, fatigue, headache, myalgia, chills, arthralgia and pyrexia.</p> <p>There have been very rare reports of myocarditis and pericarditis occurring after vaccination with Comirnaty® often in younger men and shortly after the second dose of the vaccine. These are typically mild cases and individuals tend to recover within a short time following standard treatment and rest. Healthcare professionals should be alert to the signs and symptoms of myocarditis and pericarditis. Vaccinated individuals should also seek immediate medical attention should they experience new onset of chest pain, shortness of breath, palpitations or arrhythmias.</p> <p>Individuals should be provided with the advice within the leaflet What to expect after your COVID-19 vaccination, which covers the reporting of adverse reactions and their management, such as with analgesic and/or antipyretic medication.</p> <p>Vaccinated individuals should be advised that the COVID-19 vaccine may cause a mild fever, which usually resolves within 48 hours. This is</p>

Identification and management of adverse reactions (continued)	<p>a common, expected reaction and isolation is not required unless COVID-19 is suspected.</p> <p>A detailed list of adverse reactions is available in the product's SPC.</p>
Reporting procedure of adverse reactions	<p>Healthcare professionals and individuals/carers should report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Coronavirus Yellow Card reporting scheme or search for MHRA Yellow Card in the Google Play or Apple App Store.</p> <p>As a new vaccine product, MHRA has a specific interest in the reporting of all adverse drug reactions for this product.</p> <p>Any adverse reaction to a vaccine should also be documented in the individual's record and the individual's GP should be informed.</p> <p>The Green Book Chapter 14a and Chapter 8 provide further details regarding the clinical features of reactions to be reported as 'anaphylaxis'. Allergic reactions that do not include the clinical features of anaphylaxis should be reported as 'allergic reaction'.</p>
Written information to be given to patient or carer	<p>Ensure the individual has been provided appropriate written information such as the:</p> <ul style="list-style-type: none"> • Patient Information Leaflet (PIL) for Comirnaty® COVID-19 mRNA vaccine • COVID-19 Vaccination Record Card • What to expect after your COVID-19 vaccination • COVID-19 vaccination: women of childbearing age, currently pregnant, or breastfeeding
Patient advice / follow up treatment	<p>Vaccine recipients should be monitored for 15 mins after vaccination, with a longer observation period when indicated after clinical assessment (see Chapter 14a).</p> <p>Inform the individual/carer of possible side effects and their management.</p> <p>The individual/carer should be advised to seek appropriate advice from a healthcare professional in the event of an adverse reaction.</p> <p>Vaccinated individuals should be advised to seek immediate medical attention should they experience new onset of chest pain, shortness of breath, palpitations or arrhythmias.</p> <p>Advise the individual/carer that they can report side effects directly via the national reporting system run by the MHRA known as the Coronavirus Yellow Card reporting scheme or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, they can help provide more information on the safety of medicines.</p> <p>As with all vaccines, immunisation may not result in protection in all individuals. Immunosuppressed individuals should be advised that they may not make a full immune response to the vaccine. Nationally recommended protective measures should still be followed.</p> <p>When applicable, advise the individual/carer when to return for vaccination or when a subsequent vaccine dose is due.</p>
Special considerations / additional information Continued over page	<p>Ensure there is immediate access to an anaphylaxis pack including adrenaline (epinephrine) 1 in 1,000 injection and easy access to a telephone at the time of vaccination.</p>

Special considerations / additional information
(continued)

Minor illnesses without fever or systemic upset are not valid reasons to postpone vaccination. If an individual is acutely unwell, vaccination should be postponed until they have fully recovered. This is to avoid confusing the differential diagnosis of any acute illness (including COVID-19) by wrongly attributing any signs or symptoms to the adverse effects of the vaccine.

Pregnancy

Vaccination in pregnancy should be offered in accordance with recommendations in [Chapter 14a](#), following a discussion of the risks and benefits of vaccination with the woman. Although clinical trials on the use of COVID-19 vaccines during pregnancy are not advanced, the available data do not indicate any harm to pregnancy. JCVI has therefore advised that women who are pregnant should be offered vaccination at the same time as non-pregnant women, based on their age and clinical risk group. There is extensive post-marketing experience of the use of the Pfizer BioNTech and Moderna vaccines in the USA with no safety signals so far. Over 50,000 women now report having been vaccinated whilst pregnant or when they might be pregnant in England. Because of wider experience with mRNA vaccines, these are currently the preferred vaccines to offer to pregnant women.

Routine questioning about last menstrual period and/or pregnancy testing is not required before offering the vaccine. Women who are planning pregnancy or in the immediate postpartum should be vaccinated with a suitable product for their age and clinical risk group.

If a woman finds out she is pregnant after she has started a course of vaccine, she should complete vaccination during pregnancy using the same vaccine product (unless contra-indicated).

Breastfeeding

There is no known risk associated with being given a non-live vaccine whilst breastfeeding. JCVI advises that breastfeeding women may be offered any suitable COVID-19 vaccine.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for immunisation against COVID-19; at the same time, women should be informed about the emerging safety data for the vaccine in breastfeeding. Breastfeeding women may be vaccinated under this PGD.

Previous incomplete vaccination

If the course is interrupted or delayed, it should be resumed using the same vaccine but the earlier doses should not be repeated. Evidence suggests that those who receive mixed schedules, including mRNA and adenovirus vectored vaccines make a good immune response, although rates of side effects at the second dose are higher. Therefore, every effort should be made to determine which vaccine the individual received and to complete the course with the same vaccine. For individuals who started the schedule and who attend for vaccination at a site where the same vaccine is not available or considered suitable, or if the first product received is unknown, it is reasonable to offer one dose of the locally available product to complete the primary schedule. This option is preferred if the individual is likely to be at immediate high risk or is considered unlikely to attend again. In these circumstances, this PGD may be used.

Individuals who experience severe expected reactions after a first dose of AstraZeneca or Pfizer COVID-19 vaccines appear to have a higher rate of such reactions when they receive a second dose of the alternate vaccine. Therefore, individuals who have received a first dose of the

Continued over page

Special considerations / additional information
(continued)

AstraZeneca vaccine should complete the primary course with the same vaccine, with the exception of those who experience anaphylaxis or an episode of thrombosis combined with thrombocytopaenia.

For individuals with a history of thrombosis combined with thrombocytopenia following vaccination with the AstraZeneca COVID-19 vaccine, current evidence would support completion of the course with an mRNA vaccine, provided a period of at least 12 weeks has elapsed since the dose of AstraZeneca vaccine.

For individuals with a history of capillary leak syndrome following vaccination with the AstraZeneca COVID-19 vaccine, the course may be completed with an alternative vaccine, such as Comirnaty® COVID-19 mRNA vaccine.

Individuals who are participating in a clinical trial of COVID-19 vaccines who present for vaccination should be referred back to the investigators. Eligible persons who are enrolled in vaccine trials should then be provided with written advice on whether and when they should be safely vaccinated in the routine programme.

Individuals who have been vaccinated abroad are likely to have received an mRNA or vector vaccine based on the spike protein, or an inactivated whole viral vaccine. Specific advice on completing vaccination in these individuals is available from Public Health England in the [COVID-19 vaccination: information for healthcare practitioners](#).

Co-administration with other vaccines

Where individuals in an eligible cohort present having recently received one or more inactivated or live vaccines, COVID-19 vaccination should still be given. The same applies for most other live and inactivated vaccines where COVID-19 vaccination has been received first or where an individual presents requiring two or more vaccines. It is generally better for vaccination to proceed and may be provided under this PGD, to avoid any further delay in protection and to avoid the risk of the individual not returning for a later appointment. The only exceptions to this are the shingles vaccines, where a seven-day interval should ideally be observed. This is based on the potential for an inflammatory response to COVID-19 vaccine to interfere with the response to the live virus in the older population and because of the potential difficulty of attributing systemic side effects to the newer adjuvanted shingles vaccine.

A UK study of co-administration of AstraZeneca and Pfizer BioNTech COVID-19 vaccines with inactivated influenza vaccines confirmed acceptable immunogenicity and reactogenicity. Where co-administration does occur, individuals should be informed about the likely timing of potential adverse events relating to each vaccine. If the vaccines are not given together, they can be administered at any interval, although separating the vaccines by a day or two will avoid confusion over systemic side effects.

Non-responders / immunosuppressed

Immunological response may be lower in immunocompromised individuals, but they should still be vaccinated.

JCVI advises that a third primary vaccine dose be offered to individuals aged 12 years and over who had severe immunosuppression in proximity to their first or second COVID-19 doses in the primary schedule (see 'Box: Criteria for a third primary dose of COVID-19 vaccine' in [Chapter 14a](#)). Most individuals whose immunosuppression commenced at least two weeks after the second dose of vaccination do not require an additional primary vaccination at this stage.

Continued over page

<p>Special considerations / additional information (continued)</p>	<p>The decision on the timing of the third primary dose should be undertaken by the specialist involved in the care of the individual. In general, vaccines administered during periods of minimum immunosuppression (where possible) are more likely to generate better immune responses.</p>
<p>Records</p>	<p>Record:</p> <ul style="list-style-type: none"> • that valid informed consent was given or a decision to vaccinate made in the individual's best interests in accordance with the Mental Capacity Act 2005 • name of individual, address, date of birth and GP with whom the individual is registered (or record where an individual is not registered with a GP) • name of immuniser • name and brand of vaccine • date of administration • dose, form and route of administration of vaccine • quantity administered • batch number and expiry date • anatomical site of vaccination • advice given, including advice given if excluded or declines vaccination • details of any adverse drug reactions and actions taken • supplied via PGD <p>All records should be clear, legible and contemporaneous.</p> <p>As a variety of COVID-19 vaccines are available, it is especially important that the exact brand of vaccine, batch number and site at which each vaccine is given is accurately recorded in the individual's records.</p> <p>It is important that vaccinations are recorded in a timely manner on appropriate health care records for the individual. Systems should be in place to ensure this information is returned to the individual's general practice record in a timely manner to allow clinical follow up and to avoid duplicate vaccination.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes.</p>

6. Key references

Key references	<p>Comirnaty® COVID-19 mRNA vaccine</p> <ul style="list-style-type: none">• Immunisation Against Infectious Disease: The Green Book, Chapter 14a. Updated 16 September 2021. https://www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book• Summary of Product Characteristics and Patient Information Leaflet for Comirnaty® COVID-19 mRNA vaccine. 13 August 2021 https://www.medicines.org.uk/emc/product/12740• COVID-19 vaccination programme. Updated 17 September 2021. https://www.gov.uk/government/collections/covid-19-vaccination-programme• Training recommendations for COVID-19 vaccinators. Published 8 December 2020. https://www.gov.uk/government/publications/covid-19-vaccinator-training-recommendations/training-recommendations-for-covid-19-vaccinators• National COVID-19 vaccination e-learning programme https://www.e-lfh.org.uk/programmes/covid-19-vaccination/• COVID-19 vaccinator competency assessment tool. Updated 16 March 2021 https://www.gov.uk/government/publications/covid-19-vaccinator-competency-assessment-tool• COVID-19: vaccination programme guidance for healthcare practitioners. Updated 6 August 2021. https://www.gov.uk/government/publications/covid-19-vaccination-programme-guidance-for-healthcare-practitioners <p>General</p> <ul style="list-style-type: none">• Health Technical Memorandum 07-01: Safe Management of Healthcare Waste. Department of Health 20 March 2013 https://www.gov.uk/government/publications/guidance-on-the-safe-management-of-healthcare-waste• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017. https://www.nice.org.uk/guidance/mpg2• NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017. https://www.nice.org.uk/guidance/mpg2/resources• Patient Group Directions: who can use them. Medicines and Healthcare products Regulatory Agency. 4 December 2017. https://www.gov.uk/government/publications/patient-group-directions-pgds/patient-group-directions-who-can-use-them• UK Statutory Instrument 2012 No. 1916, The Human Medicines Regulations 2012 https://www.legislation.gov.uk/ukSI/2012/1916/contents• UK Statutory Instrument 2020 No. 1125, The Human Medicines (Coronavirus and Influenza) (Amendment) Regulations 2020 https://www.legislation.gov.uk/ukSI/2020/1125/contents/made• UK Statutory Instrument 2020 No. 1594, The Human Medicines (Coronavirus and Influenza) (Amendment) Regulations 2020 https://www.legislation.gov.uk/ukSI/2020/1594/regulation/4/made
-----------------------	--

