

This patient group direction (PGD) must only be used by registered health professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used.

Patient Group Direction (PGD)

For the administration of

Heparin 5,000 units intravenous injection

For use by Ambulance Service only

Adjunctive therapy with Tenecteplase when treating ST elevation myocardial infarction (STEMI)

Throughout the Manx Care and those contracted by the Manx Care where appropriate within practice

PGD NUMBER 129

1. Change history

Version number	Change details	Date
1	Original PGD ratified	March 2022

Reference number: PGD 129 Valid from: 03/2022

Review date: 03/2024

Version: 1 Page 1 of 6

2. Medicines practice guideline 2: Patient group directions

Refer to the relevant sections of NICE medicines practice guideline 2: *Patient group directions* as stated in the blank template notes. For further information about PGD signatories, see the NHS and Manx Care <u>PGD website FAQs</u>

3. PGD development

Refer to the NICE PGD competency framework for people developing PGDs

Job Title & organisation	Name	Signature	Date
Author of the PGD			
Member of the PGD working group			

4. PGD authorisation

Refer to the <u>NICE PGD competency framework for people authorising PGDs</u>

Job Title	Name	Signature	Date
Medical Director			
Chief Pharmacist/ Pharmaceutical Adviser			
Senior Paramedic			
Director of Nursing			
GP Adviser			
Senior Microbiologist (if PGD contains antimicrobials)			

Reference number: PGD 129 Valid from: 03/2022 Review date: 03/2024

5. PGD adoption by the provider

Refer to the <u>NICE PGD competency framework for people authorising PGDs</u>

Job title and organisation	Signature	Date	Applicable or not applicable to area

6. Training and competency of registered healthcare professionals, employed or contracted by the Manx Care, GP practice or Hospice

Refer to the <u>NICE PGD competency framework for health professionals using PGDs</u>

	Requirements of registered Healthcare professionals working under the PGD	
Qualifications and professional registration	 Registered healthcare professionals, working within or contracted by the Manx Care, GP practice or Hospice who are permitted staff groups outlined within the current PGD policy Pharmacists must be practising in Manx Care authorised premises i.e. contracted pharmacy premises 	
Initial training	 Knowledge of current guidelines and the administration of the drug specified in this PGD/BNF and of the inclusion and exclusion criteria Training which enables the practitioner to make a clinical assessment to establish the need for the medication covered by this PGD Local training in the use of PGDs Successful completion of Isle of Man Ambulance Service ECG and thrombolysis training 	
Competency	Staff will be assessed on their knowledge of drugs and clinical	
assessment	assessment as part the competency framework for registered health professionals using PGDs	
Ongoing training and	The registered health care professionals should make sure they are	
competency	aware of any changes to the recommendations for this medication; it is the responsibility of the registered health care professionals to keep up to date with continuing professional development. PGD updates will be held every two years	

Reference number: PGD 129 Valid from: 03/2022 Review date: 03/2024

7. Clinical Conditions

Adjunctive therapy with Tenecteplase when treating ST elevation myocardial infarction (STEMI) PGD applies Inclusion criteria Patients aged 18 years and over Confirmed ST elevation myocardial infarction where heparin is required as adjunctive therapy with tenecteplase to reduce the risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Patient should be referred to a more experienced clinical practitioner for further assessment
Inclusion criteria Patients aged 18 years and over Confirmed ST elevation myocardial infarction where heparin is required as adjunctive therapy with tenecteplase to reduce the risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Patient should be referred to a more experienced clinical
Inclusion criteria Patients aged 18 years and over Confirmed ST elevation myocardial infarction where heparin is required as adjunctive therapy with tenecteplase to reduce the risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patient sunder 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
Confirmed ST elevation myocardial infarction where heparin is required as adjunctive therapy with tenecteplase to reduce the risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
required as adjunctive therapy with tenecteplase to reduce the risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
risk of re-infarction Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
Patient meets criteria for thrombolysis on the ambulance service thrombolysis checklist Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
thrombolysis checklist Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
Patients under 18 years of age Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
Patient does not meet the criteria for thrombolysis on the ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Patient should be referred to a more experienced clinical
ambulance service thrombolysis checklist Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Arrangements for referral Remophilia and other haemorrhagic disorders Namorrhagic disorders Necent surgery haemorrhage Severe hypertension Patient should be referred to a more experienced clinical
 Where tenecteplase is not being used as the thrombolytic drug Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
 Haemophilia and other haemorrhagic disorders Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Thrombocytopenia Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
 Recent cerebral haemorrhage Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
 Severe hypertension Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
 Severe liver disease Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
 Oesophageal varices Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Peptic ulcer Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Major trauma Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Recent surgery to eye or nervous system Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Acute bacterial endocarditis Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Elderly Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
 Spinal or epidural anaesthesia Cautions (including any relevant action to be taken) Elderly Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
Cautions (including any relevant action to be taken) • Elderly • Risk of bleeding • Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
relevant action to be taken) Risk of bleeding Severe hypertension Patient should be referred to a more experienced clinical
taken) • Severe hypertension Arrangements for referral Patient should be referred to a more experienced clinical
Arrangements for referral Patient should be referred to a more experienced clinical
·
for medical advice Intractitioner for further assessment
p
Action to be taken if Patient should be referred to a more experienced clinical
patient excluded practitioner for further assessment
• A verbal explanation should be given to the patient on: the need
patient declines for the medication and any possible effects or potential risks
treatment which may occur as a result of refusing treatment
 This information must be documented in the patients' health records
Any patient who declines care must have demonstrated capacity
to do so
Where appropriate care should be escalated

Reference number: PGD 129 Valid from: 03/2022 Review date: 03/2024

8. Details of the medicine

Name, form and strength	Unfractionated Heparin containing 5,000 units per ml.
of medicine	
Legal category	Prescription Only Medicine (POM)
Indicate any off-label use	None
(if relevant)	
Route/method of	Intravenous single bolus
administration	
Dose and frequency	Once only
	Weight <67 kg:
	Dose 4,000 units/ml
	Volume 0.8ml
	Weight ≥67 kg:
	Dose 5,000 units/ml
	Volume 1ml
Quantity to be	As per dosage table
administered	
Maximum or minimum	One dose
treatment period	
Storage	Room temperature
Adverse effects	Haemorrhage – major or minor
	Heparin induced thrombocytopenia
	Skin reactions
Records to be kept	The administration of any medication given under a PGD must be
	recorded within the patient's medical records

9. Patient information

Verbal/Written information to be given to patient or carer	 Verbal information must be given to patients and or carers for all medication being administered under a PGD Where medication is being supplied under a PGD, written patient information leaflet must also be supplied A patient information leaflet is available on request
Follow-up advice to be	If symptoms do not improve or worsen or you become unwell, seek
given to patient or carer	medical advice immediately

Reference number: PGD 129 Valid from: 03/2022 Review date: 03/2024

10. Appendix A

References

- 1. British National Formulary (BNF) available online: https://bnf.nice.org.uk
- 2. Nursing and Midwifery "The code" available online: https://www.nmc.org.uk
- 3. Current Health Care Professions Council standards of practice
- 4. Joint Royal Colleges Ambulance Liaison Committee (JRCALC) Clinical Practice Guidelines
- 5. General Pharmaceutical Council standards
- 6. The General Optical Council
- 7. Electronic medicines compendium available online: https://www.medicines.org.uk

11. Appendix B

Health professionals agreed to practice

- Each registered healthcare professional will hold their own Competency framework which will be signed and agreed by their mentor
- A mentor is defined within the Manx Care policy as any ward/area managers, sisters, senior nurses, GPs, pharmacists or senior paramedics who has completed the PGD training themselves

Reference number: PGD 129 Valid from: 03/2022 Review date: 03/2024