

Clinical Area Preparation Record – Casirivimab **600mg** and Imdevimab **600mg** in 250mL Sodium Chloride 0.9% Infusion Bag (Total volume = 260mL)
For Intravenous Infusion - Prepared Using Casirivimab and Imdevimab 300mg in 2.5mL Vials

Set up	Preparation of Infusion Bag								
<p>Step 1 Check that the dose prescribed matches the kits and preparation record selected e.g. Casirivimab 600mg and Imdevimab 600mg for Intravenous Infusion</p>	<p>Step 1 Bring 1 x Sodium Chloride 0.9% 250mL Infusion Bag from the left side of the preparation area into the middle, swab the bung with a sterile 70% alcohol wipe and allow to dry.</p>								
<p>Step 2 Casirivimab and Imdevimab infusion preparation method is complex and requires 2 persons to prepare safely:</p> <ul style="list-style-type: none"> The preparer who will prepare the product. The checker who will ensure the method is followed accurately and perform documented independent checks at key points in the preparation process. 	<p>Step 2 Bring 2 x Casirivimab 300mg in 2.5mL (120mg/mL) vials from the left side of the preparation area into the middle, swab the bung with a sterile 70% alcohol wipe and allow to dry.</p>								
<p>Step 3 Remove a kit from the refrigerator (<i>Casirivimab 600mg and Imdevimab 600mg 2.5mL vial size kit for ward preparation of Intravenous Infusion</i>) containing:</p> <ul style="list-style-type: none"> 2 cartons of Casirivimab and Imdevimab 120mg/mL Concentrate for Solution for Infusion <p>Each carton contains</p> <ul style="list-style-type: none"> 1 x Casirivimab 300mg in 2.5mL (120mg/mL) vial 1 x Imdevimab 300mg in 2.5mL (120mg/mL) vial <p>Select a Casirivimab 600mg and Imdevimab 600mg 2.5mL vial size room temperature consumable kit for ward preparation of Intravenous Infusion containing:</p> <ul style="list-style-type: none"> 1 x Sodium Chloride 0.9% 250mL Infusion Bag Syringes, needles, and administration filter 	<p>Step 3 Attach a needle to a 5mL luer lock syringe and draw up 1 x 5mL of Casirivimab 300mg in 2.5mL (120mg/mL) from the 2 vials NB: Each vial contains excess volume and therefore some liquid may remain after the required volume has been withdrawn.</p> <p>Drug name, strength and volume withdrawn checked by <i>checker (initial)</i> </p>								
<p>Step 4 Visually inspect both the Casirivimab and Imdevimab vials.</p> <ul style="list-style-type: none"> The solution for each vial should be clear to slightly opalescent, colourless to pale yellow. <p>Should particulate matter and discoloration be observed, the vial must be discarded and replaced with a new vial.</p>	<p>Step 4 Add 5mL of Casirivimab 120mg/mL to the Infusion Bag. Discard the syringe and needle into a yellow lidded sharps bin</p> <p>Addition of Casirivimab to Infusion Bag checked by <i>checker (initial)</i> </p>								
<p>Step 5 Place to the left side of the preparation area:</p> <ul style="list-style-type: none"> 2 x Casirivimab 300mg in 2.5mL (120mg/mL) vials 2 x Imdevimab 300mg in 2.5mL (120mg/mL) vials 1 x Sodium Chloride 0.9% 250ml Infusion Bag 	<p>Step 5 Place the used Casirivimab vials to the right side of the preparation area.</p>								
<p>Step 6 Prepare an infusion additive label with the following details:</p> <ul style="list-style-type: none"> Casirivimab 600mg and Imdevimab 600mg in Sodium Chloride 0.9% (Total volume = 260mL) Date and time prepared 	<p>Step 6 Bring 2 x Imdevimab 300mg in 2.5mL (120mg/mL) vials from the left side of the preparation area into the middle, swab the bung with a sterile 70% alcohol wipe and allow to dry.</p>								
	<p>Step 7 Attach a needle to a 5mL luer lock syringe and draw up 1 x 5mL of Imdevimab 300mg in 2.5mL (120mg/mL) from the 2 vials NB: Each vial contains excess volume and therefore some liquid may remain after the required volume has been withdrawn.</p> <p>Drug name, strength and volume checked by <i>checker (initial)</i> </p>								
	<p>Step 8 Add 5mL of Imdevimab 120mg/mL to the same Infusion Bag. Discard the syringe and needle into a yellow lidded sharps bin</p> <p>Addition of Imdevimab to Infusion Bag checked by <i>checker (initial)</i> </p>								
	<p>Step 9 Place the used Imdevimab vials to the right side of the preparation area</p>								
	<p>Step 10 Gently mix the filled infusion bag by slowly inverting 10 times. Do not shake</p>								
	<p>Step 11 Attach the pre-prepared label to the bag</p>								
	<p>Step 12 Ensure the product is administered using a 0.2µm to 5µm polyethersulfone, polysulfone, or polyamide end filter.</p>								
	<p>Step 13 Record details of the patient who will receive the bag below, and file the completed <i>Clinical Area Preparation Record</i> in accordance with local guidance</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">Patient Name</td> <td style="width: 25%;"></td> <td style="width: 25%;">Hospital No</td> <td style="width: 25%;"></td> </tr> <tr> <td></td> <td></td> <td>Date of Birth</td> <td></td> </tr> </table>	Patient Name		Hospital No				Date of Birth	
Patient Name		Hospital No							
		Date of Birth							