

This patient group direction (PGD) must only be used by registered health professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used.

Patient Group Direction (PGD)

For the supply of

Beclometasone Nasal Spray

By registered health care professionals for

Management of Mild Allergic Rhinitis related to Hay Fever

Throughout the Manx Care and those contracted by the Manx Care where appropriate within practice

PGD NUMBER 88

1. Change history

Version number	Change details	Date
1	Original PGD ratified	June 2021
2	Minor word changes to ensure clarity to the PGD	September 2023

Reference number: 88

Valid from: 06/2023 Review date: 09/2025 Version: 2 Page **1** of **5**

2. Medicines practice guideline 2: Patient group directions

Refer to the relevant sections of NICE medicines practice guideline 2: *Patient group directions* as stated in the blank template notes. For further information about PGD signatories, see the NHS and Manx Care <u>PGD website FAQs</u>

3. PGD development

Refer to the <u>NICE PGD competency framework for people developing PGDs</u>

Job Title & organisation	Name	Signature	Date
Author of the PGD			
Member of the PGD working group			

4. PGD authorisation

Refer to the <u>NICE PGD competency framework for people authorising PGDs</u>

Job Title	Name	Signature	Date
Medical Director			
Deputy to Chief Pharmacist/ Pharmaceutical Adviser			
Senior Paramedic			
Director of Nursing			
GP Adviser			
Senior Microbiologist (if PGD contains antimicrobials)	N/A	N/A	N/A

Reference number: 88

Valid from: 06/2023 Review date: 09/2025 Version: 2 Page **2** of **5**

5. Training and competency of registered healthcare professionals, employed or contracted by the Manx Care, GP practice or Hospice

Refer to the NICE PGD competency framework for health professionals using PGDs

	Requirements of registered Healthcare professionals working under the PGD	
Qualifications and professional registration	 Registered healthcare professionals, working within or contracted by the Manx Care, GP practice or Hospice who are permitted staff groups outlined within the current PGD policy Pharmacists must be practising in Manx Care authorised premises i.e. contracted pharmacy premises 	
Initial training	 Knowledge of current guidelines and the administration of the drug specified in this PGD/BNF and of the inclusion and exclusion criteria Training which enables the practitioner to make a clinical assessment to establish the need for the medication covered by this PGD Local training in the use of PGDs 	
Competency	Staff will be assessed on their knowledge of drugs and clinical	
assessment	assessment as part the competency framework for registered health professionals using PGDs	
Ongoing training and	The registered health care professionals should make sure they are	
competency	aware of any changes to the recommendations for this medication;	
	it is the responsibility of the registered health care professionals to	
	keep up to date with continuing professional development. PGD	
	updates will be held every two years	

6. Clinical Conditions

Clinical condition or	Seasonal mild rhinitis related to hay fever	
situation to which this		
PGD applies		
Inclusion criteria	Adults and children over 6 years presenting with symptoms of	
	seasonal mild rhinitis related to hay fever	
Exclusion criteria	Under 6 years	
	Recent nasal surgery or untreated nasal infection	
	Pulmonary tuberculosis	
	 Any known hypersensitivity to any component of the 	
	medicine	
	Patient refuses treatment under this Patient Group Direction	

Reference number: 88

Valid from: 06/2023 Review date: 09/2025 Version: 2 Page **3** of **5**

Cautions (including any	Avoid after nasal surgery (until healing has occurred)
relevant action to be	Avoid in pulmonary tuberculosis
taken)	Avoid in the presence of untreated nasal infections
	Patients transferred from systemic corticosteroids may
	experience exacerbation of some symptoms
	Systemic absorption may follow nasal administration particularly
	if high doses are used or if treatment is prolonged; therefore
	also consider the cautions and side-effects of systemic
	corticosteroids
	A detailed list of cautions is available in the SPC, which is available
	from the electronic Medicines Compendium website:
	www.medicines.org.uk and BNF https://bnf.nice.org.uk
Arrangements for referral	Patient should be referred to a more experienced clinical
for medical advice	practitioner for further assessment
Action to be taken if	Patient should be referred to a more experienced clinical
patient excluded	practitioner for further assessment
Action to be taken if	A verbal explanation should be given to the patient on: the need
patient declines	for the medication and any possible effects or potential risks
treatment	which may occur as a result of refusing treatment
	This information must be documented in the patients' health
	records
	Any patient who declines care must have demonstrated capacity
	to do so
	Where appropriate care should be escalated

7. Details of the medicine

Name, form and strength	Beclometasone Nasal Spray (50micrograms/metered Spray)
of medicine	
Legal category	Prescription Only Medicine (POM)
Indicate any <u>off-label use</u>	None
(if relevant)	
Route/method of	Intranasal
administration	
Dose and frequency	100 micrograms (2 sprays) into each nostril twice daily, reduced to
	50 micrograms (1 spray) into each nostril twice daily when
	symptoms controlled
Quantity to be supplied	1 x 200 dose metered spray
Maximum or minimum	Maximum: Total 400micrograms (8 sprays) daily
treatment period	
Storage	Room temperature
Adverse effects	Altered smell sensation
	Epistaxis
(continued)	Headache
	Nasal complaints
	Taste altered
	Throat irritation
	Glaucoma

Reference number: 88
Valid from: 06/2023 Review date: 09/2025 Version: 2 Page **4** of **5**

Adverse effects	Nasal septum perforation (more common following nasal surgery)
(continued)	 Vision blurred. MHRA/CSM advice: patients should be advised to report any blurred vision or other visual disturbances – rare risk of central serous chorioretinopathy Systemic absorption may follow nasal administration particularly if high doses are used or if treatment is prolonged. Therefore also consider the side-effects of systemic corticosteroids
	A detailed list of adverse reactions is available in the SPC, which is available from the electronic Medicines Compendium website:
	<u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u>
Records to be kept	The administration of any medication given under a PGD must be
	recorded within the patient's medical records

8. Patient information

Verbal/Written information to be given to patient or carer	 Verbal information must be given to patients and or carers for all medication being administered under a PGD Where medication is being supplied under a PGD, written patient information leaflet must also be supplied A patient information leaflet is available on request
Follow-up advice to be	If symptoms do not improve or worsen or you become unwell, seek
given to patient or carer	medical advice immediately

9. Appendix A

References

- 1. British National Formulary (BNF) available online: https://bnf.nice.org.uk
- 2. Nursing and Midwifery "The code" available online: https://www.nmc.org.uk
- 3. Current Health Care Professions Council standards of practice
- 4. General Pharmaceutical Council standards
- 5. Electronic medicines compendium available online: https://www.medicines.org.uk

10. Appendix B

Health professionals agreed to practice

- Each registered healthcare professional will hold their own Competency framework which will be signed and agreed by their mentor
- A mentor is defined within the Manx Care policy as any ward/area managers, sisters, senior nurses, GPs, pharmacists or senior paramedics who has completed the PGD training themselves

Reference number: 88

Valid from: 06/2023 Review date: 09/2025 Version: 2 Page **5** of **5**